

**M. Sc. (MATHEMATICS WITH
APPLICATIONS IN COMPUTER
SCIENCE) [M. Sc. (MACS)]**

Term-End Examination

December, 2024

MMT-006 : FUNCTIONAL ANALYSIS

Time : 2 Hours

Maximum Marks : 50

Weightage : 70%

Note : (i) *Question No. 6 is compulsory.*

(ii) *Attempt any **four** of the remaining questions.*

1. (a) Is the map $A : \mathbf{R}^3 \rightarrow \mathbf{R}^2$, $A(x, y, z) = (x + y, y + z)$ open ? Justify. 2

(b) Show that the operator A on l^2 defined by

$$Ax(n) = \frac{i^n}{n} x(n) \text{ is compact and normal. } 2+2$$

(c) Prove that c_0 is a closed subspace of c and

$$\dim \frac{c}{c_0} = 1. \quad 2+2$$

2. (a) Show that on a Hilbert space, Hahn-Banach extensions are unique. 2

- (b) For $x \in l^1$, define $f(x) = \sum \frac{n}{n+1} x(n)$. Prove that f is a bounded linear functional with $\|f\| = 1$ and there is no $x \neq 0$ with $|f(x)| = \|x\|_1$. 1+1+2

- (c) Let M be a closed subspace of a normed space X . Show that $d(x, M) = \|x\|$ if and only if there is a bounded linear functional $f \neq 0$ on X such that $f = 0$ on M and $|f(x)| = \|f\| \|x\|$. 2+2

3. (a) Define $\langle \cdot, \cdot \rangle : \mathbb{C}^4 \times \mathbb{C}^4 \rightarrow \mathbb{C}$ as :

$$\langle x, y \rangle = x(1) \overline{y(1)} + x(2) \overline{y(2)} + x(3) \overline{y(3)} - x(4) \overline{y(4)}$$

Show that $\langle \cdot, \cdot \rangle$ is linear in the 1st variable and conjugate symmetric but not an inner product. 3+3

- (b) Define A on $C[0, 1]$ by $Af(t) = t f(t)$. Prove that A is a bounded operator, but is not compact. 2+2

4. (a) State bounded inverse theorem. Use the theorem to show the following :

Suppose H is a Hilbert space and A is a bounded linear operator such that $\langle Ax, x \rangle \geq 0$ for all $x \in H$. Assume that A defines a complete norm on H given by $\|x\|_A^2 = \langle Ax, x \rangle$.

Show that $\exists c$ such that $\langle Ax, x \rangle \geq c \|x\|^2$.

1+2

- (b) Let X be a Banach space and let $c_0(X)$ be the space of all sequences (x_n) in X such that $\lim \|x_n\| = 0$. Prove that $c_0(X)$ is a Banach space with norm $\|(x_n)\| = \sup \|x_n\|$.

3

- (c) If M is a closed subspace of a Hilbert space H , determine $M^{\perp\perp}, M^{\perp\perp\perp}, \dots$.

4

5. (a) Let H be a Hilbert space $u \in H, u \neq 0$. Define $Ax = \langle x, u \rangle u, x \in H$ and

$Bx = \langle x, u \rangle \frac{u}{\|u\|}$. Calculate A^*, A^2 and B^2 .

3

- (b) Let $X = C_{00}$, $u_n = (0, 0, \dots, 0, 1, 0, 0, \dots)$, where only the n th entry is 1 and let $k_n = \frac{1}{n}$ for $n = 1, 2, \dots$. Show that Riesz-Fischer theorem does not hold good. 3
- (c) Define $A : l^p \rightarrow l^p$ by $Ax = (0, x_1, x_2, \dots)$, $1 \leq p < \infty$. Show that A is an isometry which has no eigen values. What is $R(A)$ in terms of $\{e_n\}$? 3+1
6. State, with justification, whether the following statements are True or False : 5×2=10
- (a) If a subspace M of a normed space X is finite dimensional, then $\dim \frac{X'}{M^\perp} < \infty$.
- (b) A Banach space cannot be the union of countably many proper closed subspaces.
- (c) If A is a bounded operator on l^2 and $A^2 = 0$, then $A = 0$.
- (d) There is no linear map $C^3 \rightarrow C^4$ that takes any orthonormal basis to an orthonormal set.
- (e) There is a linear map $R^m \rightarrow R^n$ with a closed graph that is not continuous.

× × × × × × ×