

No. of Printed Pages : 8

MCH-013

**M. SC. (CHEMISTRY)/M. SC. IN
ANALYTICAL CHEMISTRY
(MSCCHEM AND MSCANCHEM)**

Term-End Examination

June, 2025

MCH-013 : GENERAL PHYSICAL CHEMISTRY

Time : 2 Hours

Maximum Marks : 50

Note : *Question No. 1 is compulsory. Attempt any **four** questions from question nos. 2 to 7.*

Use the following values whenever it requires :

$$k_B = 1.38 \times 10^{-23} \text{ JK}^{-1}; h = 6.626 \times 10^{-34} \text{ J-s};$$

$$N_A = 6.022 \times 10^{23} \text{ mol}^{-1}, R = 8.314 \text{ JK}^{-1}$$

$$\text{mol}^{-1}; 1 \text{ cm}^{-1} = 1.98 \times 10^{-23} \text{ J}.$$

1. Attempt any *five* questions from the following : 5×2=10

- (i) Outline the advantages of the stopped flow method over continuous flow methods.
- (ii) Write any *two* limitations of the Debye-Hückel's law. In which type of solution this law holds good ?
- (iii) What is the mechanism of enzyme action which involves the binding of the substrates ?
- (iv) Write any *two* limitations of Lindemann's theory of unimolecular reaction.
- (v) What are amorphous materials ? Give any *two* examples.
- (vi) What is the difference between isobaric and isochoric processes ?

- (vii) Write Stirling's approximation and outline its limitations.
- (viii) Define the flux in transport phenomenon and outline its significance.
2. (a) Write the expression for drift speed in terms of viscosity η and name the terms involved. 2
- (b) Can a catalyst alter the equilibrium constant for a reaction ? Justify your answer. 2
- (c) Calculate the mean activity coefficient, γ_1 of 0.001 M aqueous solution of a Na_2SO_4 at 25°C . 3
- (d) Using Stirling's approximation, calculate the % error introduced in computing the value of $6!$ [Use $\ln 2 = 0.7$ and $\ln 3 = 1.1$]. 3

3. (a) Define Fick's second law of diffusion and outline its significance in terms of curvature of the concentration gradient. 3
- (b) The relaxation time for a given fast equilibrium reaction $A \xrightleftharpoons[k_{-1}]{k_1} B$ is found to be $5 \mu\text{s}$ from temperature jump method. If the equilibrium constant is 1.4×10^{-3} , calculate the rate constant for forward and reverse reactions. 3
- (c) For an isothermal and isobaric process the change in entropy and enthalpy are found to be 175 J/K and 54 kJ/mol , respectively. Calculate the following : 4
- (i) Whether this process is spontaneous at 57°C ?

- (ii) At which temperature the system will attain equilibrium, assuming that the enthalpy and entropy changes are not affected by the change in the temperature ?
4. (a) In a binary mixture of an ideal gas in which $n_A = 7.5n_B$. If the partial molar volume of B is decreased by $0.05 \text{ m}^3/\text{mol}$, what would be the change in partial molar value of A ? 2
- (b) Write the mathematical expression of most probable molecular speed, u_{mp} and name the terms involved. 2
- (c) Calculate the angle (in $\sin^{-1} \theta$) at which the second order reflection will occur when X-rays of wavelength 0.16 nm are diffracted by the atom of a crystal, given that the inter-planar distance is 0.4 nm . 2


- (d) Derive the relationship between the term activation energy in Arrhenius equation and the critical barrier energy proposed in collision theory. 4
5. (a) Write the rate determining step under high pressure condition for Lindemann unimolecular reaction. 2
- (b) Why the pressure-jump method is better for studying the gaseous system ? Give reasons. 2
- (c) Calculate the relative population of two energy levels separated by 1000 cm^{-1} at 27°C .

Or

Calculate the rotational partition function at 500 K for a diatomic molecule having moment of inertia of $2.70 \times 10^{-43}\text{ kgm}^2$. 3

- (d) For Michaelis-Menten mechanism for enzyme substrate system, the reaction rate changes from first order to the zero order. Give the physical reason for this. 3
6. (a) Write the physical significance of turnover number in Michaelis-Menten mechanism. 2
- (b) Draw a labelled schematic diagram showing the potential energy profiles of catalyzed and uncatalyzed reactions. 2
- (c) What are the basic assumptions of the Debye-Hückel's theory ? Why do we use water as an electrolyte in this model ? 3
- (d) At room temperature, sodium crystallizes in a body centered cubic cell with $a = 0.4$ nm. Calculate the density of sodium. [Given : Molar mass of sodium = 23.0 g/mol.] 3

7. (a) Calculate the standard Gibbs energy change for the following reaction at 300 K :



Given that :

5

	N_2O	H_2	N_2H_4	H_2O
$\Delta H^\circ_f (\text{kJ/mol})$	80	0	50	– 240
$S^\circ (\text{J/K})$	220	130	120	190

- (b) Derive the translational partition function for a particle of mass m confined into a one-dimensional box of length l , where the energy of the particle is given as :

5

$$\epsilon_{n, \text{trans}} = \frac{n^2 h^2}{8ml^2}$$

× × × × ×